

Synergy® Dry Pasture 16:10

Liquid Supplement #124427

(For Ruminants Only)

Fat-fortified, protein supplement for beef cattle.

NOT FOR HUMAN CONSUMPTION

Guaranteed Analysis

Crude Protein	Min.	16.0 %
(This includes not more than 10.0% equivalent crude protein from non-protein nitrogen.)		
Crude Fat	Min.	10.0 %
Crude Fiber	Max.	0.1 %
Acid Detergent Fiber (ADF)	Max.	0.1 %
Phosphorus (P)	Min.	0.7 %
Dry Matter.....	Min.	48.0 %
Moisture	Max.	52.0 %

Ingredients

Condensed Fermented Corn Extractives, Condensed Extracted Glutamic Acid Fermentation Product, Condensed Soybean Solubles, Vegetable Oil, Lecithin, Salt, Urea, Sulfuric Acid, Xanthan Gum.

CAUTION: USE ONLY AS DIRECTED.

Use only as recommended on the reverse side.

SUGGESTION: USE OF THIS PRODUCT MAY RAISE THE MOISTURE CONTENT OF FEEDS TO WHICH IT IS APPLIED. IF ANTICIPATED FINAL FEED MOISTURE IS TO EXCEED 12% AND STORAGE FOR PERIODS LONGER THAN 2 TO 3 DAYS IS EXPECTED, SPECIAL CONSIDERATION TO THE INCLUSION OF PRESERVATIVES IS NEEDED. YOUR WESTWAY REPRESENTATIVE WILL PROVIDE ASSISTANCE IF NEEDED.

SELLER MAKES NO WARRANTY, EXPRESSED OR IMPLIED, AS TO DESCRIPTION, UTILITY, PRODUCTIVENESS, OR ANY OTHER MATTER, EXCEPT AS TO GUARANTEED ANALYSIS.

This product was manufactured in a facility that does not handle or store animal protein products prohibited in ruminant feed as described in 21 CFR section 589.2000 and 589.2001.

This feed was produced in a facility certified in the American Feed Industry Association's Safe Feed/Safe Food Certification Program; for details go to: www.safefeedsafefood.org.

Manufactured by:



Westway Feed Products LLC

14015 Park Drive, Suite 217
Tomball, TX 77377



Liquid Bulk Shipment



Daily recirculation is required, whether in use or not. Product must be stored in a recirculation tank, see instructions below:

- Liquid supplements stored in re-circulating tank systems: Re-circulate immediately prior to use moving no less than 1 percent of the tank contents per minute from the bottom of the tank to the top for not less than 10 minutes. Re-circulate daily, as described above, even when supplement is not used.
- Liquid supplements stored in mechanical, air or other agitation-type tank systems: Agitate immediately prior to use for not less than 10 minutes, creating a turbulence at the bottom of the tank that is visible at the top. Agitate daily, as described even when supplement is not used.

1. For Dairy and Beef cattle of all classes this product can be fed as a part of grain mixes or total mixed rations at a rate not to exceed the provision of 1 pound of crude protein equivalent per 1000 pounds of animal body weight.

2. For free choice lick wheel feeding, the following precautions must be observed:

- This product contains supplemental selenium and therefore cannot be feed free choice to lactating dairy cattle.
- Provide unlimited drinking water.
- This product is a supplement. Adequate roughage, both in quantity and quality, must be available at all times.
- Grain should be fed as necessary.
- Animals must have consumed other feed prior to being given this Liquid Feed Supplement for the first time. Do not provide overly hungry animals free access to this Liquid Feed Supplement.
- If animals are not fed a Liquid Feed Supplement for 48 hours, observe animals closely for three days to prevent overconsumption.
- Provide a free choice mineral supplement containing adequate levels of calcium, phosphorus, salt, and trace minerals.
- Lick tanks must be in good working order. Loose or damaged tank lids, open lick wheel holes, and leaks must be repaired before animals are allowed access to lick tanks

It is not recommended that this product be fed free choice or by lick wheel feeders. If customer decides to feed in this manner, it is done so at their own risk. Positional stability of this product is not warranted.